



2017 Cow-Calf Protocol

Modified Live Virus (MLV) Vaccine Protocol:

Open Cows: Vaccinated 4 to 8 weeks prior to breeding

Bovi-Shield GOLD FP 5 VL5 (Optional Hb combo if needed)

Ultra Choice 7

Dectomax Injectable or Dectomax Pour On

Calves 2 to 3 Months of Age—Vaccinated at the same time cows are given pre-breeding vaccines:

Inforce 3 and One Shot BVD or Bovi-Shield GOLD One Shot

UltraChoice 7

Dectomax Injectable or Dectomax Pour On

*If banding include a tetanus toxoid vaccine according to mfg. directions

Killed BVD Virus Vaccine Protocol:

Bred Cows:

CattleMaster GOLD FP 5 L5

UltraChoice 7

Dectomax Injectable or Dectomax Pour On

Calves 2 to 3 Months of Age:

CattleMaster FP 5

One Shot

UltraChoice 7

Dectomax Injectable or Dectomax Pour On

*If banding include a tetanus toxoid vaccine according to mfg. directions

*Work with your veterinarian for their recommendations.



Modified Live Virus (MLV) Vaccine Protocol:

Option 1:

Weaning:

Inforce 3 and One Shot BVD
UltraChoice 7
Dectomax Injectable or Dectomax Pour On

Post Weaning (14-21 days after weaning):

Bovi-Shield Gold 5
UltraChoice 7

Option 2:

Weaning:

Bovi-Shield GOLD One Shot
UltraChoice 7
Dectomax Injectable or Dectomax Pour On

Post Weaning (14-21 days after weaning):

Bovi-Shield GOLD 5
UltraChoice 7

Killed Virus Vaccine Protocol:

Option 3:

Pre-Weaning working (2-4 weeks prior to weaning):

CattleMaster FP 5
Ultrabac 7 Somubac
Dectomax Injectable or Dectomax Pour On

Weaning:

Bovi-Shield GOLD One Shot
Ultrabac 7 Somubac