



# 2017 Tennessee Beef Alliance Health Protocol



Select one brand, one box, & an Approved Wormer for a complete protocol

**All HEIFERS must have a shot of Lutalyse or SYNCHSURE on second working!**



### Pre-Weaning

#### 1<sup>st</sup> Working (2 weeks prior to weaning)

- TRIANGLE 5
- Presponse HM
- Bar-Vac 7/Somnus

\*Use Approved Wormer\*

#### 2<sup>nd</sup> Working (14-28 days later) (Following Weaning)

- EXPRESS 5 or Pyramid 5
- Bar-Vac 7/Somnus

### Post or at Weaning

#### 1<sup>st</sup> Working

- EXPRESS 5 HS
- Presponse HM

\*Use Approved Wormer\*

#### 2<sup>nd</sup> Working (14-28 days later)

- Express 5 HS
- Alpha-7 or Alpha-7/MB-1

### Post or at Weaning

#### 1<sup>st</sup> Working

- Pyramid 5 + Presponse SQ
- Bar-Vac 7/Somnus

\*Use Approved Wormer\*

#### 2<sup>nd</sup> Working (14-28 days later)

- Pyramid 5
- Bar-Vac 7/Somnus

### Pre-Weaning

#### 1<sup>st</sup> Working (2 weeks prior to weaning)

- Cattlemaster GOLD FP 5
- Ultrabac 7 Somubac

\*Use Approved Wormer\*

#### 2<sup>nd</sup> Working (14-28 days later) (Following Weaning)

- Bovi-Shield GOLD ONE SHOT (Combo)
- Ultrabac 7 Somubac

### Post or at Weaning

#### 1<sup>st</sup> working

- Bovi-Shield GOLD ONE SHOT (combo)
- Ultrabac 7 Somubac

\*Use Approved Wormer\*

#### 2<sup>nd</sup> Working (14-28 days later)

- Bovi-Shield GOLD 5
- Ultrabac 7 Somubac

### \*Approved Wormers\*

Wormers listed are approved to be used with either program. **NO GENERIC WORMERS**

- Cydectin Injectable or Pour-on
- Dectomax Injectable or Pour-On
- Eprinex
- Ivomec Injectable or Pour-on
- Long Range

**For More Information on Health Protocols contact:**

Dr. C. F. Locklar – Vet. Advisor  
931-388-4750

Dr. Tiffany Krantz-Vet. Advisor  
931-403-5553

Dr. George Boyd-Vet. Advisor  
931-528-8805

Stacey Turner – Boehringer  
615-585-5337

Todd Jennings – Boehringer  
931-842-0450

Barry Maupin - Zoetis  
731-487-2877

**PLEASE MAINTAIN ALL SERIAL AND LOT NUMBERS FROM VACCINES FOR HEALTH RECORDS**

\*Note if over 100 days past last shot, another round of 2<sup>nd</sup> working shots must be given.

\* 2<sup>nd</sup> shot completed no later than 2 weeks prior to sale, or more than 100 days prior to sale.